



**GLYCYRRHIZA GLABRA L. O'SIMLIGINING KIMYOVIY TARKIBI
ХИМИЧЕСКИЙ СОСТАВ РАСТЕНИЯ GLYCYRRHIZA GLABRA L.
CHEMICAL COMPOSITION OF GLYCYRRHIZA GLABRA L. PLANT**

Mustaqil izlanuvchi: **Xoldarov I.I**

<https://doi.org/10.5281/zenodo.6642730>

Annotatsiya

Risolada Glycyrrhiza glabra L. o'simligining botanik tavsifi, tarqalishi hamda kimyoviy tarkibi yoritib o'tilgan.

Kalit so'zlar: Glycyrrhiza glabra L, Fabaceae, shirimiya, triterpen saponin, glisterrizin kislota, flavonoid, xalkon, isoflavon, efir moylari va uchuvchan moddalar.

Аннотация

В статье описано ботанические характеристика, распространение и химический состав растения Glycyrrhiza glabra L.

Ключевые слова: Glycyrrhiza glabra L, Fabaceae, солодка, тритерпеновый сапонин, глицерризиновая кислота, флавоноид, халкон, изофлавон, эфирные масла и летучие вещества.

Annotation

The article describes the botanical characteristics, distribution and chemical composition of the plant Glycyrrhiza glabra L.


Key words: Glycyrrhiza glabra L, Fabaceae, licorice, triterpene saponin, glisterrhizic acid, flavonoid, chalcone, isoflavone, essential oils and volatiles.

Shirinmiya (Glycyrrhiza glabra L.) –dukkakdoshlar (Fabaceae) oilasiga mansub ko'p yillik o't o'simlik. Glycyrrhiza glabra L. foydali dorivor o'simliklardan biridir. Glycyrrhiza qadimgi yunoncha glykos atamasidan kelib chiqqan bo'lib, shirin va rhiza atamasi ildiz degan ma'noni anglatadi. Glycyrrhiza glabra L. Shimoliy Hindistonda mulaithi nomi bilan mashhur. Glycyrrhiza glabra L., shuningdek, qizilmiya va shirin yog'och deb ham ataladi, vatani O'rta er dengizi va Osiyoning ayrim hududlari [1,112].


Teofrast o'z asarlarida bu o'simlikni solodkoviy koren, skifskaya trava, pontiyskaya trava nomi bilan atagan. Mazkur o'simlikni rus tilida – solodka golaya, o'zbek tilida-shirinmiya, chuchukmiya, qizilmiya, Qoraqalpog'iston respublikasida esa bo'yan deb atashadi. Glycyrrhiza glabra vatani Evroosiyo, Shimoliy Afrika va G'arbiy Osiyo. U Afrikada (Liviya); Osiyo (Armaniston, Ozarbayjon, Gruziya, Rossiya Federatsiyasi, Xitoy, Qozog'iston, Qirg'iziston, Tojikiston,



Turkmaniston, O'zbekiston, Mo'g'uliston, Eron, Iroq, Afg'oniston, Falastin, Jordaniya, Livan, Suriya, Turkiya, Hindiston, Pokiston); Evropa va Rossiya Federatsiyasi - Evropa qismida tarqalgan [2,12]. Respublikamiz hududida shirinmiya tipik to'qay o'simligi hisoblanib, asosan Sirdaryo va Amudaryoning quyi qismlarida uchraydi. Shirinmiya polikarp o't o'simlik, poyalari yaxshi rivojlangan bo'lib, silindrsimon tuzilishga ega. Poya yog'ochlangan bo'lib balandligi 150-160sm, ba'zan uning balandligi to'qay sharoitlarda 200 sm va undan ham ortadi. Sho'rangan tuproq sharoitida bu ko'rsatkichlar 50-70sm atrofida qayd etiladi.



Shirimmiya ildizi triterpen saponinlardan (4-20%), asosan glitserrizin, ya'ni 18β -glisterzin kislotani kaliyli va kalsiyli tuzlari (o'simlikni yer usti qismi asosan birikmasi bo'lib, glitserrizin kislotasi yoki glitserrizin saqlab, shakarga nisbatan 50 marta shirindir. Shirinmiya ildizi liquiritic kislota, glycyrrhetic acid, glycyrrhizic acid, isoglabrolide, and liquorice kislota kabi kislotalarni saqalydi. 18β -glycyrrhetic acid ($3-O-(2-O-\beta-d\text{-glucopyranuronosyl}-\alpha-d\text{glucopyranosyl})-3-\beta\text{-hydroxy}-11\text{-oxo}-18\beta,20\beta\text{-olean-12-en-29-oic acid}$) Glycyrrhiza glabra ildizi dan ajratib olingandir[3,171].





Nº	Fitobirikmalar	Tajribani o'tkazilishi	Natija
1	Uglevodlar	Molish sinovi	(-)
2	Oqsillar	Mis sulfat sinovi	(-)
3	Flavonoidlar	Qo'rg'oshin asetat sinovi, NaOH eritmasi sinovi	(+)
4	Alkaloidlar	Dragendorf sinovi	(+)
5	Steroidlar	Liberman sinovi	(+)
6	Terpenoidlar	Salkovskiy sinovi	(+)
7	Saponinlar	Ko'pik sinovi	(+)
8	Tanninlar	Temir(III) xlorid sinovi	(+)
9	Flobatanninlar	HCl sinovi	(-)
10	Antraxinonlar	Benzol sinovi	(-)
11	Glikozidlar	Keller-Kiliani sinovi	(+)
12	Fenol birikmalar	Temir sulfat sinovi	(-)

Glycyrrhiza glabra o'simligidan quidagi flavonoid va chalconlar ajartib olingan: liquiritin, liquiritigenin, hamnoliquiritin, neoliquiritin, isoliquiritin, isoliquiritigenin, neoisoliquiritin, licuraside, glabrolide, licoflavonol, 5,8-dihydroxy-flavone-7-O-beta-D-glucuronide, glychionide A



va 5-hydroxy-8-methoxylflavone-7-O-beta-D-glucuronide va glychionide B. Flavonoidlar shirinmiyaning sariq rangiga javob beradi.

Glycyrrhiza glabra o'simligidan quidagi isoflavonlar ajratib olingan: glabridin, galbrene, glabrone, shinpterocarpin, licoisoflavone A and B, formononetin, glyzarin, kumatakenin, hispaglabridin A, hispaglabridin B, 4'-O-methylglabridin and 3'-hydroxy-4'-O-methylglabridin, glabroisoflavanone A vad B glabroiso-flavanone B [4,678].

Glycyrrhiza glabra barglaridan gidrodistillatsiya usuli bilan ajratib olingan efir moylari GC va GC-MS yordamida o'rganilib, quyidagi asosiy uglevodorod va kislorod tutgan birikmalar aniqlangan: isoniazid (13.36%), diethyltoluamide (6.56 %), benzoic kislota (5.37 %), benzene (4.58 %), linalool (2.25 %), prasterone (5.63 %), warfarin (1.43 %), iodoquinol (1.90 %), phenol, 4-(2-aminopropyl)(1.30 %). Glycyrrhiza glabra ildizidan ajratib olingan efir moylarida 82 birikma borligi aniqlanib, asosiy birikmalari quyidagilardir: hexanoic kislota (31.57%), hexadecanoic kislota (3.30%), hexanol (1.71%) va octanoic kislota(1.44%). Efir moylarining hidi estragole (methyl chavicol), anethole, eugenol, indole, γ -nonalactone va cumic spirtlarini mavjudligi bilan bog'liqdir [5,1120]. Misr, Afg'oniston, Suriya, Xitoy va Germaniyada o'suvchi namunalar uchuvchan moddalarning tarkibi va miqdori bilan bir-biridan farq qiladi. Glycyrrhiza glabra o'simligining ildizidan ajratib olingan uchuvchan moddalari tarkibida quyidagi birikmalar aniqlangan: (E)-2-heptenal, 5-methyl-furfural, (2E, 1E) heptadienol, (E)-2-octen-1-al, o-guaiacol, 2-phenylethanol, (Z)-pinene hydrate, lavandulol, terpinen-4-ol, (E)-linalool oxide, p-cymen-8-ol, α -terpineol, methyl chavicol, (4E)-decenal, decanal, (2E, 4E)-nonadienal, cumin aldehyde, carvone, piperitone, (E)-cinnamaldehyde, (E)-anethole, (2E, 4Z)-decadienal, thymol, indole, carvacrol, (2E, 4Z)-decadienal, p-vinylguaiacol, eugenol, γ -nonalactone, methyl eugenol, β -caryophyllene, β -dihydro-ionone, himachalene epoxide, spathulenol, (1 α , 10 α)-Epoxy-amorph- 4-ene, β -caryophyllene oxide va humulene epoxide II[6,525].

Amalda qo'llanilayotgan TIF TN kod raqamlari

TIF TN kodlari

Pozitsiyaga izoh



2938

2938 10 000 0

2938 90

2938 90 100 0

2938 90 300 0

2938 90 900 0

2938 90 900 1

2938 90 900 2

2938 90 900 3

Tabiiy yoki sintetik glikozidlар, ularning tuzлari, oddiy va murakkab efirlари va boshqa hosilalari

- rutin va uning hosilalari
- boshqalar:
- tirnoqgul glikozidlari
- glisirrizin kislotasi va glisirrizinatlar
- boshqalar:
- qizilmiya suvli ekstrakti
- qizilmiya metanolli ekstrakti
- qizilmiya etanolli ekstrakti

Foydalanilgan adabiyotlar

1.J.-M. Mérillon, K.G. Ramawat (eds.), Sweeteners, Reference Series in Phytochemistry, 2018,pp.112.

2.Ali Esmail Al-Snafi " Glycyrrhiza glabra: A phytochemical and pharmacological review" IOSR Journal of Pharmacy (IOSRPHR), Vol. 8, №06.2018, pp. 01-17.

3.Isbrucker RA and Burdock GA. Risk and safety assessment on the consumption of Licorice root (Glycyrrhiza sp.), its extract and powder as a food ingredient, with emphasis on the pharmacology and toxicology of glycyrrhizin. Regular Toxicol Pharmacol 2006; 46: 167-192.

4. Li JR, Wang YQ and Deng ZZ. Two new compounds from Glycyrrhiza glabra. J Asian Nat Prod Res 2005; 7: 677-680.

5.Kameoka H and Nakai K. Components of essential oil from the root of Glycyrrhiza glabra. Nippon Nageikagaku Kaishi 1987; 61(9): 1119-1121.

6. Quirós-Sauceda AE, Ovando-Martínez M, Velderrain-Rodríguez GR, González-Aguilar GA and AyalaZavala JF. Licorice (Glycyrrhiza glabra Linn.) oils. In: Essential Oils in Food Preservation, Flavor and Safety, Edited by Preedy VR. First Edition 2016; Chapter 60: 523-530.